



## **Effects of Aqueous Extract of *Ocimum gratissimum* Leaves and Vitamin C on Lead Acetate-induced Changes in the Thymus of Adult Wistar Rats**

**G. N. Okechukwu<sup>1\*</sup>, E. Ezor<sup>1</sup>, E. Finbarrs-Bello<sup>2</sup>, L. N. Ebube<sup>1</sup>, G. C. Uzomba<sup>1</sup> and A. O. Ibegbu<sup>1</sup>**

<sup>1</sup>*Department of Anatomy, Faculty of Basic Medical Sciences, Federal University Ndufu-Alike Ikwo (FUNAI), Ebonyi State, Nigeria.*

<sup>2</sup>*Department of Anatomy, Faculty of Basic Medical Sciences, Enugu State University (ESUT), Parklane, Enugu State, Nigeria.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author GNO designed the study, author GCU performed the statistical analysis, wrote the protocol and Author EFB wrote the first draft of the manuscript. Authors AOI and LNE managed the analyses of the study. Author EE managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJBCRR/2019/v26i130087

#### Editor(s):

(1) Dr. Halit Demir, Professor, Department of Chemistry, Faculty of Art and Science Yuzuncu, Yil University, Turkey.

#### Reviewers:

(1) Senthil Kumar Raju, Swamy Vivekanandha College of Pharmacy, Tiruchengode, Tamilnadu, India.

(2) S. Murugesan, Pachaiyappa's College, University of Madras, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49098>

**Original Research Article**

**Received 27 February 2019**

**Accepted 14 May 2019**

**Published 04 June 2019**

### **ABSTRACT**

Lead is a heavy metal found in earth's crust. It is a widespread and insidious environmental toxin known as a severe and aggressive contaminant to human and animal organisms' health status. This work is aimed at evaluating the effect of aqueous extract of *Ocimum gratissimum* (OG) on lead induced changes in the thymus of adult albino wistar rats. Thirty five male Wistar rats were used in the study and were randomly divided into seven groups with five rats in each group. The rats in Group one (G1) served as the Control and received distilled water. Group 2 received 120 mg/kg body weight (bwt) of lead acetate, G3 received 375 mg/kg bwt of OG only. G4 received 120 mg/kg bwt of lead acetate and OG extract at 375 mg/kg bwt., G5 received 120 mg/kg bwt of lead and OG at 750 mg/kg bwt. G6 received 375 mg/kg bwt of OG in two weeks followed by 120 mg/kg bwt of

\*Corresponding author: E-mail: okeygetrude@gmail.com;

lead acetate for one week, while G7 received 120 mg/kg bwt of lead acetate and Vitamin C at 11900 mg/kg. All the administrations were carried out orally for twenty one (21) days. At the end of the administration, the rats were fasted for 24 hours. They were weighed and humanely sacrificed via cervical dislocation. The thymus was harvested in all the groups and prepared for histological studies using routine haematoxylin and eosin (H&E) staining techniques. The result of the present study shows that lead possessed a distortive effect in the histoarchitecture of thymus of the wistar rat in G2 when compared with G1. Treatment of these experimental animals with vitamin C proved to possess more ameliorative effect in restoring the histoarchitecture on lead toxicity in the thymus closer to normal than OG. Therefore, it is suggested that aqueous extract OG can act via the same pathway as vitamin C, in maintaining the normal histological structures of the thymus of adult albino wistar rat exposed to lead toxicity.

**Keywords:** *Ocimum gratissimum*; thymus; Vitamin C; lead.

## 1. INTRODUCTION

Various harmful and toxic chemical compounds (lead and mercury) are formed as by/intermediate products of normal biochemical reactions in the human system; these by products are eliminated or detoxified under normal physiological conditions. Lead is a heavy metal found in earth's crust. Free radicals and reactive oxygen species (ROS) are amongst these chemicals and their accumulation when not eliminated by the endogenous system often lead to oxidative stress [1]. Lead is a chemical element with symbol Pb (from the Latin plumbum) and atomic number 82. It is a heavy metal that is denser than most common materials. Lead is soft and malleable, and has a relatively low melting point. When freshly cut, lead is bluish-white; it tarnishes to a dull gray color when exposed to air. Lead has the highest atomic number of any stable element and concludes three major decay chains of heavier elements. Lead is not naturally present in the human body and, as it currently exists in the environment, has been identified as a health threat [2]. As suggested by various authors, [3]; [4] lead may contaminate humans from various pathways including: inhalation of airborne lead particulates, consumption of water or food contaminated by lead, and ingestion (due to contamination of hands or other objects) of soil or dust contaminated with lead. Once in the body, lead travels in the blood to soft tissues such as the liver, kidneys, lungs, brain, spleen, muscles, and heart [3].

Cells are equipped with different kinds of mechanisms to fight against radicals and reactive oxygen species and to maintain redox homeostasis of the cell. For example, antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx)

play important roles in scavenging the free radicals and preventing injury [5]. The antioxidant might play a role in the treatment of lead poisoning [6]. Vitamins like C is capable to control the critical balance of oxidants and antioxidants in the body. [7] and [8] reported that intake of ascorbic acid decrease the blood lead level and control the lead toxicity.

*Ocimum gratissimum* belongs to the group of plants known as spices. The plant is an erect small plumb with many barnacles usually not more than 1 m high [9] and [10]. In South East Asia, it is cultivated as a home garden crop but it is grown on a commercial scale in Vietnam. In Nigeria, it is Efinrin in Yoruba, Diadoyal in Hausa and Nchuanwu in Igbo [11]. In traditional medicine, the leaves have been used as a general tonic and anti-diarrhoea agent and for the treatment of conjunctivitis by instilling directly into the eyes; the leaf oil when mixed with alcohol is applied as a lotion for skin infections, and taken internally for bronchitis [12]. Although, conventional antibiotics have been very useful in orthodox medicine, it has been argued by many that its concomitant use with herbal extracts is not desirable as one normally antagonizes the activity of the other [13].

The thymus gland is the main organ of the lymphatic system. Its primary function is to promote the development of specific cells of the immune system called T-lymphocytes. Once mature, these cells leave the thymus and are transported via blood vessels to the lymph nodes and spleen. T-lymphocytes are responsible for cell-mediated immunity, which is an immune response that involves the activation of certain immune cells to fight infection.

In addition to immune function, the thymus also produces hormones that promote growth and maturation [14].

## 2. MATERIALS AND METHODS

### 2.1 Collection and Identification of Plant Materials

The fresh leaf of *Occimum gratissimum* was obtained behind the Department of Anatomy Federal University, Ndufu-Alike, Ikwo. The specimen was identified by Botanist at Department of Biology Alex-Ekwueme Federal University, Ndufu-Alike, Ikwo with the specimen number 1285. The leaves were washed and shade dried.

### 2.2 Experimental Animals

In order to minimize variations in response, only adult male rats were used for this study. Thirty-five adult albino wistar rats, weighing between 140-160 g were obtained from the Animal House of the Department of Human Anatomy, Alex-Ekwueme Federal University Ndufu-Alike Abakili, Ebonyi State, Nigeria. They were maintained under favorable condition at standard temperature and pressure and fed with top food grower mash (Vital Feed®). Prior to acclimatization which lasted for two weeks, the rats were divided into seven groups, each of five rats. They were fed with feed and water *ad libitum* throughout the experiment.

### 2.3 Plant Extraction Procedure

It was extracted in the Department of chemistry, University of Nigeria, Nsukka. The dried specimen were grounded into powder of 500 g using a Q-link electric blender; Model QBL-18L40, and stored in air-tight containers [15]. Aqueous infusion was done by mixing a calculated volume of distilled water and powdered sample. The mixture was allowed to stand for 30 minutes before filtration. It was then centrifuged at about 3000xg for 5 min and the supernatant collected

### 2.4 Experimental Design

The design consist of thirty-five (35) animals, randomly divided into seven (7) groups each of five rats in each group, and the rats were fed with animal feed and water *ad libitum*. Distilled water were used to dissolve the *Occimum gratissimum* extract, the lead acetate and vitamin C. The treatment was done orally for 21 days. After the administration (24 hours), the rats were sacrificed and the organs harvested. The rats in Group one (G1) served as the Control and received distilled water. Group 2 received 120

mg/kg body weight (bwt) of lead acetate, G3 received 375 mg/kg bwt of OG only [16,17]. G4 received 120 mg/kg bwt of lead acetate and OG extract at 375 mg/kg bwt., G5 received 120 mg/kg bwt of lead and OG at 750 mg/kg bwt. G6 received 375 mg/kg bwt of OG in two weeks followed by 120 mg/kg bwt of lead acetate for one week, while G7 received 120 mg/kg bwt of lead acetate and Vitamin C at 11900 mg/kg [18].

### 2.5 Animal Sacrifice

The animals were sacrificed by cervical dislocation followed by the midline incision to harvest the organ.

The thymus was procured and cleared from adjoining supporting, connective tissues. The organ was gently blotted with filter paper and weighed. The thymus was immediately immersed in 10% formalin solution to be processed for haematoxylin and eosin staining and light microscopic studies.

### 2.6 Tissue Preparation for Histology

Following sacrifice, the tissue was prefixed in 10% formal saline, after whole body perfusions were transferred to a graded series of ethanol. On Day 1, they were placed in 70% alcohol for 7 hours, then transferred to 90% alcohol and left in the latter overnight. On day 2, the tissues were passed through three changes of absolute alcohol for an hour each and then cleared in xylene. Once cleared, the tissues were infiltrated with molten paraffin wax in the oven at 58°C. Three changes of molten paraffin wax at one-hour interval was made, after which the tissues were embedded in wax and blocked out. Prior to embedding, it was ensured that the mounted sections to be cut by the rotary microtome were orientated perpendicular to the long axes. These sections were designated "vertical sections". Serial sections of 4µm thick were obtained from a solid block of tissue, fixed on clean slides to which Mayer's egg albumin had been coated to cement the sections to the slides properly and were stained. For the histological and histochemical study, sections were stained with H & E, after which they were passed through a mixture of equal concentration of xylene and alcohol. Following clearance in xylene the sections were oven-dried between 35°C and 40°C [19,20]. The sections were examined under the light microscope at high power magnification (x 100 and x 400) and photomicrographs taken.

## 2.7 Statistical Analysis

Statistical analysis was done using both electronic calculate and SPSS version 20. All data were expressed as mean  $\pm$  SD of a number of experiments (n = 5). The values will be compared using the Student T-test. P value less than 0.05 (P < 0.05) was considered as being significant.

## 3. RESULTS

It was observed that there was a gradual increase in the mean body weight of the animals in group 1 (control group), and other groups of experimental animals except the animals treated

with lead only (Groups 2) and the animals treated with high dose of *Ocimum grattissimum* and lead (group 5) which shows a decrease in the mean body weight.

Thymus exposed to lead show a significant decreased in normal thymic lymphocytes and reticular epithelial cells compared to control. Thymic atrophy was observed in all the groups administered with lead acetate but the occurrence was severe in animal treated with lead acetate only (group 2). This atrophy was slight in animals treated with lead acetate and OG, (group 4, 5 and 6) and moderate in animal treated with lead acetate combined with vitamin C (group 7).





Fig. 1.

## 3.1 Body Weight Difference


Table 1. Weight change and weight comparison of Wistar rats at Initial (day 1) and Final (day 21) day of treatment

Groups	Initial weight (gram)	Final weight	ange in weight (gram)	Percentage change in weight (%)
1	89.90 $\pm$ 15.93	114.5 $\pm$ 12.99	24.6 $\pm$ 2.94*	27.36
2	159.1 $\pm$ 25.06	153.9 $\pm$ 42.84	-5.2 $\pm$ 17.78**	3.27
3	127.4 $\pm$ 33.84	151.9 $\pm$ 49.99	24.5 $\pm$ 16.15*	19.23
4	108.7 $\pm$ 24.00	137.7 $\pm$ 19.34	29 $\pm$ 4.46*	26.68
5	201.4 $\pm$ 20.27	189.1 $\pm$ 33.11	-12.3 $\pm$ 12.84**	6.11
6	110.5 $\pm$ 6.816	151.2 $\pm$ 24.99	40.7 $\pm$ 18.17*	36.83
7	118.7 $\pm$ 15.23	145.0 $\pm$ 11.79	26.3 $\pm$ 3.44*	22.17


The values are expressed as Mean  $\pm$  STD \*significant increase \*\*significant decrease. (P < 0.05)  
 Weight change and weight comparison of Wistar rats at Initial (day 1) and Final (day 21) day of treatment



**Fig. 2.** Photomicrograph of thymus sections from (group 1) , with rat thymus showing; CTL- cortex with thymic lymphocytes, TB- trabecular, BV- blood vessels, ERC-epithelia reticular cells, and HC- hassall's corpuscles(H&E x 400)



**Fig. 3.** Photomicrographs (A and B) of thymus sections from (group 2) , with rat thymus showing; C-cortex with dense aggregation of thymic lymphocytes, TB- trabecular, M- medullar with distorted epithelia reticular cells, and HC- distorted hassall's corpuscles (H&E x 400)



**Fig. 4.** Photomicrograph of thymus sections from (group 3) showing normal and well oriented; HC-hassall's corpuscle, ERC- epithelia reticular cell, BV-blood vessels and CTL- cortex with thymic lymphocytes (H&E x 400)





Fig. 6. Photomicrograph of thymus sections from (group 5) showing; shrunken BV- blood vessel, ERC- epithelia reticular cell, distorted or damaged, HC- Hassall's corpuscle, (H&E x400)





Fig. 7. Photomicrograph of thymus sections from (group 6) showing damaged BV-blood vessels and HC- Hassall's corpuscle (H&E X 400) Fig. 5. Photomicrograph of thymus sections from (group 4) showing well oriented; BV-blood vessels, TB- trabecular, M-medulla with epithelia reticular cell and CTL- cortex with thymic lymphocytes (H&E x 400)



**Fig. 8. Photomicrograph of thymus section from (group 7) showing damaged TB-trabecular and MTL- medulla with thymic lymphocytes**

## 6. DISCUSSION

This study showed that there was reduction in mean body weight of the rats in the lead treated groups only when compared with the rats in the Control group. This observed reduction in the mean bodyweight is in agreement with [24] who stated that rats that are continuously exposed to heavy metals such as mercury (Hg), Cadmium (Cd), Arsenic (As) and Lead (Pb) usually results in reduction in the body weight. This reduction in the mean body weight with continuous exposure to lead could be explained on the basis of anorexia (loss of appetite) which is caused by lead exposure [22]. The decrease in the mean body weight could be explained more on the basis that the reduction of the mean body weight may be due to the decreased muscle mass and cachexia as a result of the oxidative stress induced by lead. In that, there was evidence that heavy metal toxicity was linked to oxidative stress [23] which according to many researchers was associated with muscle wasting and cachexia leading to low body weight [24].

Furthermore, this study agrees with [25,26,27] who postulated that generation of reactive oxygen species, stimulation of lipid peroxidation and depletion of antioxidants are the major contributors to lead exposure related biochemical and histopathological changes in the animal body organs.

More so, as seen from the animals in G4 and G5, the administration of OG at low dose is an effective therapeutic strategy in restoring the

histoarchitecture of the thymus as reported by Patrick, 2006 in one of his studies where he stated that lead exposed animals showed that a therapeutic strategy to increase antioxidant defense system of the body may be of help for long-term effective treatment of lead poisoning. Recently, several chelating agents approaches have been proposed therapeutically, including supplementation with antioxidants and up-regulation of endogenous anti-oxidative defense system for lead induced oxidative stress in various body organs [28,29,30]. However, the mechanism of actions of these chelating agents and antioxidants is still indistinct. It is believed that chelating agents reduce the lead toxicity in soft organs through its chelating activity, whereas antioxidants protect the cells from influence of oxidative damage protect the cells from influence of oxidative damage by scavenging the free radical generation and inhibiting of lipid peroxidation [31,32,33,23]. As indicated by this study and [28], chelating agent, like OG, reduces the toxic effect of heavy metals on the histological sections of the thymus (G4 and G5), although it does not have similar effect in protecting the tissues against toxic effect (G6). Finally this study is in agreement with the believed that vitamins with antioxidant activity protect soft tissues against damaging effect of free radicals which produced as a result of heavy metals toxicity [29].

## 7. CONCLUSION

It could be right to conclude that aqueous extract of *O. gratissimum* in a dose-dependent manner has been proven to be an attractive target very

effective in its protective effects against lead acetate induced thymus damages and act via the same route as vitamin C.

### ETHICAL APPROVAL

The ethical committee of the Faculty of Basic Medical Sciences Alex-Ekwueme Federal University, Ndufu-Alike, Ikwo approved the study with ethical number MPC/1707/02/003.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

- Halliwell B. How to characterize an antioxidant: An update. *Biochem. Soc. Symp.* 1995;61:73-101.
- Atlanta Georgia. Toxicological profile for cadmium. ATSDR, agence for toxic substance and disease registry; 1998.
- Watemaux C, et al. Methodological issues in modeling the relationship between low – level lead exposure and infant development: Examples from the boston lead study. *Environ Res.* 1985;38:119-129.
- Sharma RP, Street JC. Public health aspects of toxic heavy metals in animal feeds. *J Am Vet Med Assoc.* 1980;177: 149-153.
- Bergendi L, Benes L, Durackova Z, Ferenci KM. Chemistry, physiology and pathology of free radicals. *Life Science.* 1999;65:1865-1874.
- Gurer H, Ozygunes H, Saygin E, Ercal N. Antioxidant effect of *Taurine* against lead - induced oxidative stress. *Arch. Environ. Contam. Toxicol.* 2001;41:397-402.
- Simon JA. Relationship of ascorbic acid to blood leads levels. *JAMA.* 1999;281:2289-2293.
- Hudes ES. Relationship of ascorbic acid to blood leads levels. *JAMA.* 1999;281:2289-2293.
- Vierra RF. Chemical characterization of *Ocimum gratissimum* found in the market and used in Traditional medicine in Brazil. *Journal of Economic Botany.* 2000;20:5-6.
- Simon JE. Chemical characterization of *Ocimum gratissimum* found in the market and used in Traditional medicine in Brazil. *Journal of Economic Botany.* 2000;20:5-6.
- Owulade MO. Handbook African medicinal plants. CRC Press, Boca Raton, Florida, USA. 2004;214-215.
- Iwu MM. Handbook of African medicinal plants. CRC Press, New York. 1993;214-215.
- Nakamura CV, Ishid K, Faccin LC, Filho BP, Cortez DA, Rozental S, Souza W, Ueda N. *In vitro* activities of essential oils from *Ocimum gratissimum* against four *Candida* species. *Research in Microbiology.* 2004;155(7):579-86.
- Lymphoid organs; 2018. Available:<http://teachmeseries.com> (Assessed 23 June 2018)
- Odey MO, Mboso O. Nat prod leaves were blended into fine powder using electric blender model QBL -18L40. *Biotechnology Advances J. Biotech;* 2012.
- Ebeye OA, Ekundina OV, Wilkie IE. Histological and biochemical effects of aqueous extract of *Ocimum gratissimum* on the liver and kidney of adult wistar rats, *African Journal of Cellular Pathology.* 2014;2:59-64.
- Ofem OE, Ani EJ, Eno AE. Effect of aqueous leaves extract of *Ocimum gratissimum* on hematological parameters in rats. *International Journal of Applied Asic Medical Research.* 2012;2(1):38–42.
- Material Safety Data Sheet (MSDS). Safety data for mercuric chloride physical and chemical properties. Janssen Pharmaceutical 3a. 2440 Geel, Belgium; 2008.
- Sheehan DC, Hrapchak BB. Theory and practise of histotechnology. Battelle Memorial Institue; 1987.
- Eisler R. Lead hazards to fish, wildlife, and invertebrates, asynoptic review, Biological Report 85(1.14), Contaminant Hazard Reviews 14, Laurel, MD, USA: US Fish and Wildlife Service, 1988 (Last modified in 2013); 1988.
- Nwokocha CR, Owu DU, Ufearo CS, Iwuala MO. Comparative study on the efficacy of *Garcinia kola* in reducing some heavy metal accumulation in liver of Wistar rats. *Journal of Ethnopharmacology.* 2011;135:488-91.
- Klaassen CD. Heavy metals and heavy-metal antagonists In: Goodman and Gilman's Pharmacological Basis of Therapeutics. (10<sup>th</sup> edition.), McGraw-Hill, New York; 2001.
- Yeh YC, Liu TJ, Wang LC, Lee HW, Ting CT. A standardized extract of *Ginkgo biloba* suppresses doxorubicin-induced oxidative stress and p53- Mediated mitochondrial apoptosis in at testes.

- British Journal of Pharmacology. 2009; 156:48-61.
24. Buck M, Chojkier M. Muscle wasting and dedifferentiation induced by oxidative stress in a murine model of cachexia is prevented by inhibitors of nitric oxide synthesis and antioxidants. EMBO Journal. 1996;15:1753-1765.
25. Silbergeld EK, Waalkes M, Rice JM. Lead as carcinogen: Experimental evidence and mechanism of action. Am. J. Indus. Med. 2000;38:316-323.
26. Patrick L. Lead toxicity part II. The role of free radical damage and the use of antioxidants in the pathology and treatment of lead toxicity. Altern. Med. Rev. 2006;11:114-127.
27. Jabeen R, Tahir M, Raza SMH, Nisa FU, Batool S. Protective role of vitamin C against teratogenic effects of lead acetate on mouse kidney. 2011;1-5. Available:<http://pjmhsonline.com>
28. Raafat BM, El-Barbary A, Touson E, Aziz S. Di-mercapto succinic acid (DMSA) and vitamin C chelating potent in lead intoxication, regarding oxidative stress and apoptotic related proteins in rabbits. Journal of Genetic Engineering and Biotechnology. 2011;9(2):121-131.
29. Hamadouche Slimani M, Aoues A. Beneficial effect of administration of vitamin C in amelioration of lead hepatotoxicity. Nat. Sci. Biol. 2012;4(3):7-13.
30. Dewanjee S, Sahu R, Karmakar S, Gangopadhyay M. Toxic effects of lead exposure in Wistar rats: Involvement of oxidative stress and the beneficial role of edible jute (*Corchorus olitorius*) leaves. Food Chem. Toxicol. 2013;55:78-91.
31. Bashandy SA. Beneficial effect of combined administration of vitamin E in amelioration of chronic lead hepatotoxicity. Egypt J Hosp Med. 2006;23:371-84.
32. Haleagrahara T, Jackie S, Chakravarthi M, Rao M, Kulur A. Protective effect of *Etilingera eletior* (torch ginger) extract on lead acetate-induced hepatotoxicity in rats. J. Toxicol. Sci. 2010;35(5):663-671.
33. El-Neweshy MS, El-Sayed YS. Influence of vitamin C supplementation on lead-induced histopathological alterations in male rats. J. Exp. Toxicol. Pathol. 2011; 63:221-227.

© 2019 Okechukwu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/49098>