



## Toxic Mechanisms of Aryloxyphenoxypropionates in Target and Non-target Organisms

Ayokanmi Ore<sup>1\*</sup> and Ebenezer Tunde Olayinka<sup>1</sup>

<sup>1</sup>Department of Chemical Sciences, Biochemistry Division, Ajayi Crowther University, Oyo, Nigeria.

### **Authors' contributions**

This review was prepared in collaboration between both authors. Author AO prepared the first draft of the manuscript. Author ETO reviewed the manuscript. Both authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/IJBCRR/2019/v27i130109

#### Editor(s):

(1) Dr. Cheorl-Ho Kim, Professor, Department of Biological Science, Molecular and Cellular Glycobiology Unit, Sungkyunkwan University, South Korea.

#### Reviewers:

(1) Toungos, Mohammed Dahiru, Adamawa State University, Nigeria.  
(2) Zvonko Pacanoski, Ss. Cyril and Methodius University, Republic of Macedonia.  
Complete Peer review History: <http://www.sdiarticle3.com/review-history/48220>

Review Article

Received 29 March 2019  
Accepted 07 June 2019  
Published 03 August 2019

### **ABSTRACT**

Herbicides are substances used to control unwanted plants-weeds. They can be classified into several classes by mechanism of action. This review describes the members of aryloxyphenoxypropionate herbicides, their pharmacokinetic properties, metabolism and their mechanism of phytotoxicity in target weeds as well as in non-target organisms. Two major toxicity mechanisms are described. The first is by inhibition of lipid synthesis. This is achieved by inhibiting the rate limiting step of lipid biosynthesis catalyzed by acetyl CoA carboxylase. The second mechanism is by induction of oxidative stress. This is achieved by generation of reactive oxygen species which in excess can cause oxidative damage to macromolecules and cellular structures especially the membrane lipids. Loss of vital membrane lipids alters the fluidity of membrane, loss of cellular contents and eventually cell death and death of the entire plant.

**Keywords:** Herbicides; Aryloxyphenoxypropionates; toxicity mechanisms; acetyl CoA carboxylase; oxidative stress.

\*Corresponding author: E-mail: [a.ore@acu.edu.ng](mailto:a.ore@acu.edu.ng), [oreayokanmi@gmail.com](mailto:oreayokanmi@gmail.com);

## 1. INTRODUCTION

Herbicides (also known as weedkillers), are chemical substances used to control unwanted plants (weeds) [1]. Herbicides are described as either selective or non-selective. Selective herbicides are used to control specific weed species, while leaving the desired crop unharmed. The non-selective herbicides (total weedkillers) are applied to kill all plant species and therefore can be used to clear waste ground, construction sites, industrial sites etc. Apart from classification as selective/ non-selective, other classifications include persistence (by residual action), mechanism of uptake (whether they are absorbed through foliage, or through the roots, or by other means), and mechanism of phytotoxic action (how they kill susceptible plants). Herbicides may also be distinguished by application as pre-emergence or post-emergence. Pre-emergent herbicides are applied to the soil before the crop emerges and prevent germination of weed seeds and early growth of weeds. Post-emergent herbicides are applied after the crop and weeds have emerged [2].

Aryloxyphenoxypropionic herbicides or aryloxy phenoxypropionates (ArOPPs) belongs to the class of phenoxy herbicides and the sub-class of phenoxypropionic herbicides [3]. ArOPPs are relatively new class of selective herbicides prepared from heterocyclic oxyphenoxy propionic acid derivatives for the control of graminaceous weeds in broad-leaved crops and tolerant cereal crops [4]. They are applied mainly on cotton, soybeans and sugar beet as well as in the wheat, corn, barley, rye, legumes, sunflowers, sesame, alfalfa, peanuts, lettuce, spinach, potatoes, cucumber, peas, tomatoes, fennel, and strawberries [5-8]. ArOPPs are post-emergence (PoEm) herbicides composed of about fifteen members. Esters of ArOPPs are newer forms of PoEm herbicides with higher selectivity. In spite of a wide range of chemical structures, they display similar herbicidal action. The Common names, IUPAC names, Chemical structures and Chemical Abstracts Service registry numbers of AOPP herbicides are shown in Table 1.


## 2. ABSORPTION, TRANSLOCATION AND METABOLISM OF ArOPPs

Following application to fields, herbicides (their active ingredients) must be absorbed significantly for effective herbicidal actions. In phytotoxicity studies, absorption refers to the quantity of applied active ingredient that has been absorbed

by the leaves or roots of the treated plants. Distribution is the fractionation of accumulated herbicides in parts of the treated plant including treated leaf (or root) [9].








### 2.1 Absorption, Translocation and Metabolism in Target Plant Species








In plants, ArOPPs are absorbed mainly through the leaves. In few cases, they may be absorbed through the roots. Studies on the absorption of fluazifop-butyl (FB) through the leaves, for instance, was found to be about 75% after 8 hours of application [10]. Translocation of FB from leaves to root is also very rapid with up to 10 fold higher in the leaves compared to the root [11]. Similar absorption and translocation rates are also observed with other ArOPPs. A study carried out on cyhalofop-butyl showed an absorption of up to 73% in *Echinochloa* spp. (a major weed in rice) twenty-four hours after application [12]. Absorption/ translocation studies conducted by Aguero-Alvarado et al. [13] in soybean also revealed a significantly high absorption/ translocation of haloxyfop. Following absorption and translocation, they are metabolised into the acidic forms. The esters of ArOPPs undergo a more rapid decomposition, yielding the corresponding free acids as the main metabolites [14]. Fluazifop-p butyl (FPB), one of the most extensively studied ArOPPs, has been shown to be metabolized mainly to fluazifop-p (also known as fluazifop acid) and other minor metabolites (Fig. 1). The major metabolites of haloxyfop-p butyl and cyhalofop-butyl have also been found to be haloxyfop-p and cyhalofop-butyl acid respectively [13,15]. These acidic metabolites are then translocated to the entire plant through the phloem and xylem systems, accumulating in the meristematic tissues of the plant [16].








**Fig. 1. Metabolism of fluazifop-p-butyl (a) to fluazifop-p (b) and other minor metabolites [17]**

**Table 1. Common names, IUPAC names, Chemical structures and Chemical Abstracts Service (CAS) registry numbers of AOPP herbicides**

S/N	Common Names	IUPAC Nomenclature	Chemical Structures	CAS No
1	Chlorazifop	(RS)-2-[4-(3,5-dichloro-2-pyridyloxy)phenoxy]propionic acid		60074-25-1
2	Clodinafop	(R)-2-[4-(5-chloro-3-fluoro-2-pyridyloxy)phenoxy]propionic acid		114420-56-3
3	Clofop	(RS)-2-[4-(4-chlorophenoxy)phenoxy]propionic acid		26129-32-8
4	Cyhalofop	(R)-2-[4-(4-cyano-2-fluorophenoxy)phenoxy]propionic acid		122008-78-0
5	Diclofop	(RS)-2-[4-(2,4-dichlorophenoxy)phenoxy]propionic acid		40843-25-2
6(a)	Fenoxaprop	(RS)-2-[4-(6-chloro-1,3-benzoxazol-2-yloxy)phenoxy]propionic acid		95617-09-7
(b)	Fenoxaprop-P	(R)-2-[4-(6-chloro-1,3-benzoxazol-2-yloxy)phenoxy]propionic acid		113158-40-0

S/N	Common Names	IUPAC Nomenclature	Chemical Structures	CAS No
7	Fenthiaprop	(RS)-2-[4-(6-chloro-1,3-benzothiazol-2-yloxy)phenoxy]propionic acid		66441-11-0
8(a)	Fluazifop	(RS)-2-[4-[5-(trifluoromethyl)-2-pyridyloxy]phenoxy]propionic acid		69806-34-4
(b)	Fluazifop-P	(R)-2-[4-[5-(trifluoromethyl)-2-pyridyloxy]phenoxy]propionic acid		83066-88-0
9(a)	Haloxifop	(RS)-2-[4-[3-chloro-5-(trifluoromethyl)-2-pyridyloxy]phenoxy]propionic acid		69806-34-4
(b)	Haloxifop-P	(R)-2-[4-[3-chloro-5-(trifluoromethyl)-2-pyridyloxy]phenoxy]propionic acid		95977-29-0
10	Isoxapyrifop	(RS)-2-[2-[4-(3,5-dichloro-2-pyridyloxy)phenoxy]propionyl]isoxazolidine		87757-18-4
11	Kuicaoxi	ethyl(2RS)-2-[(2RS)-2-[4-(6-chloroquinoxalin-2-yloxy)phenoxy]propionyloxy]-3-methylbut-3-enoate		3724-55-8

S/N	Common Names	IUPAC Nomenclature	Chemical Structures	CAS No
12	Metamifop	(R)-2-[4-(6-chloro-1,3-benzoxazol-2-yloxy)phenoxy]-2'-fluoro-N-methylpropionanilide		256412-89-2
13	Propaquizafop	2-isopropylideneaminoxyethyl(R)-2-[4-(6-chloroquinoxalin-2-yloxy)phenoxy]propionate		111479-05-1
14	Quizalofop	(RS)-2-[4-(6-chloroquinoxalin-2-yloxy)phenoxy]propionic acid		76578-12-6
	Quizalofop-P	(R)-2-[4-(6-chloroquinoxalin-2-yloxy)phenoxy]propionic acid		94051-08-8
15	Trifop	(RS)-2-[4-(α,α,α-trifluoro-p-tolyloxy)phenoxy]propionic acid		58594-74-4

## 2.2 Absorption, Pharmacokinetics and Metabolism in Human and Animals

Absorption of ArOPPs are also very rapid in animals by oral route. In a study on rats, cyhalofop-butyl was rapidly absorbed following oral dosing with maximum plasma levels attained within a few hours. The major metabolite in plasma, liver, bile, kidney, urine and feces, was found to be cyhalofop-butyl acid (66-78%) and a minor metabolite, 4-(4-cyano-2-fluorophenoxy) phenol. Absorption of ArOPPs through the skin have also been reported in human and animal studies [18,19]. FPB is absorbed orally and through the skin, although at a slower rate through the skin [19,20].

## 3. PHYTOTOXICITY MECHANISMS OF ArOPPs

Considerable efforts have been made towards understanding the mechanism of phytotoxic action of the ArOPP herbicides. These involved studies cutting across a wide range of plant species. Major phytotoxicity mechanisms identified include inhibition of lipid metabolism and oxidative stress.


### 3.1 Inhibition of Lipid Metabolism

#### 3.1.1 Acetyl CoA carboxylase

All the known aryloxyphenoxypropionates show similar species specificity and similar symptoms in susceptible plants. Based on these similarities, it has been suggested that they have similar targets in susceptible plant species. Studies have postulated that ArOPPs acts by disabling the *de novo* fatty acid biosynthesis, but no specific site of action was identified [21,22]. Studies have however demonstrated that they specifically inhibit acetyl-CoA carboxylase (ACCase) in target plants. ACCase (E.C. 6.4.1.2) is the rate-limiting enzyme in the *de novo* fatty acid biosynthetic pathway [23]. This plastid located enzyme catalyzes the adenosine triphosphate (ATP)-dependent formation of malonyl-CoA from acetyl-CoA and bicarbonate (Fig. 2). Despite differences in regulation and protein structure across various species, ACCase is known to play the same role in fatty acid synthesis [24].

Fatty acids are the major components of the cell membrane. It is viewed that the inhibition of *de novo* fatty acid biosynthesis may lead to low production of relevant fatty acids. This leads to impaired membrane biosynthesis and loss of cellular components [26]. Several esters of ArOPP such as diclofop, fenoxaprop, fluzifop,

haloxyfop, quizalofop etc were found to cause similar effects in most plant species [26].



**Fig. 2. Acetyl CoA carboxylase catalyzed reaction; the first step in the sequence of reactions for the biosynthesis of fatty acids [25]**

#### 3.1.2 Fatty acid synthase

Fatty acid synthase (FAS) has been implicated as another site of action for the graminicides [27]. Fatty acid synthase is the enzyme system that catalyzes the synthesis of saturated long-chain fatty acids from acetyl CoA, malonyl CoA, and NADPH. Inhibition of fatty acid synthase results in a decrease in the incorporation of malonyl-coenzyme A into fatty acids, but does not affect the incorporation of acetate into malonyl-coenzyme A.

### 3.2 Oxidative Stress

Generation of oxidative stress has also been identified as one of the mechanism used by ArOPPs to kill susceptible plant species. This is achieved by generation of reactive species (RS), depletion of antioxidant defenses, and generation of oxidative stress. Oxidative stress can be defined as a serious imbalance between the production of reactive species and antioxidant defenses [28]. Reactive species are chemically reactive chemical species containing oxygen (reactive oxygen species, ROS), nitrogen (reactive nitrogen species, RNS) etc. ROSs (which are the most extensively studied RS) are oxygen-containing molecules exhibiting higher chemical reactivity than O<sub>2</sub>. ROS is constantly generated in the chloroplasts due to partial reduction of O<sub>2</sub> or as a result of transfer of energy to O<sub>2</sub>.

ROS may play two different roles in plant physiology; low levels can be used in adaptive signalling whereas, high concentrations of ROS can result in phytotoxicity [29]. The role of ROS in plant physiology has been extensively described [30]. Oxidative stress can also be defined as a disturbance in the pro-oxidant-antioxidant balance in favour of the pro-oxidants, leading to cell damage. An antioxidant is any substance that can inhibit the oxidation of the cell components such as DNA, proteins and lipids.

Damages resulting from oxidative stress are often called 'oxidative damages'. Oxidative damage to macromolecules (lipids, proteins, DNA etc) results in formation of oxidative damage products (lipid, protein, DNA oxidation products) which are usually considered biomarkers of oxidative stress [31]. Several levels of antioxidative defense mechanism are used by plants to prevent oxidative damages [32]. Table 2 shows the major reactive species, antioxidants and biomarkers of oxidative damage. Several studies have shown ArOPPs to cause oxidative stress or oxidative damage as a phytotoxicity mechanism in plant species. Studies on ArOPPs across a wide variety of plants are summarized in Table 3.

#### 4. TOXICITY MECHANISMS OF ArOPPs IN NON-TARGET ANIMAL SPECIES

Studies evaluating the toxicity of ArOPPs in non-plants has described similar toxicity mechanisms as those found in plant models. Studies in human and animal including bacterial models show inhibition of lipid metabolism and or generation of oxidative stress.

##### 4.1 Inhibition of Lipid Metabolism

Metabolic studies indicate that ArOPPs are metabolized to the acidic metabolites similar to

plant. Fluazifop-p butyl for instance is metabolised to fluazifop acid in human and animals models [18,19]. These metabolites also appear to produce similar symptoms in mammals. Also, weight loss observed in mammalian studies have been linked to inhibition of mammalian ACCase by fluazifop compounds [45]. In a study conducted by Kemal and Casida [46], they observed inhibition of rat liver ACCase activity by fluazifop-P-butyl by competitive inhibition mechanism. It was further observed that fluazifop-P and fluazifop-P-butyl may have a lesser affinity for mammalian ACCase, compared with the ACCase in sensitive plants.

##### 4.2 Oxidative Stress

Oxidative stress has also been linked to the toxicity mechanism elicited by ArOPPs in animal studies. A study carried out by Ye et al. [47] in which cyanobacteria was exposed to diclofop and its R-enantiomer reported an increase in MDA concentration and SOD activity. They suggested that diclofop and its R-enantiomer may cause a collapse in the transmembrane proton gradient and damage the cell membrane through lipid peroxidation and free radical attack. Exposure of rat to fluazifop-p butyl and haloxyfop-p methyl ester caused depletion in hepatic and testicular antioxidant defence system,

**Table 2. Major reactive oxygen species, antioxidants and biomarkers of oxidative damage in plants [32,33]**

<b>Major Reactive Oxygen Species (ROS)</b>	
Free Radicals	Hydroxyl radical ( $OH^*$ ); Superoxide radical ( $O_2^{\cdot-}$ )
Non-radicals	Singlet oxygen ( $^1O_2$ ); Hydrogen peroxide ( $H_2O_2$ )
<b>Major cellular antioxidants and their protective functions</b>	
Enzymic	Superoxide dismutase (SOD): converts $O_2^{\cdot-}$ to $H_2O$ and $O_2$ Catalase (CAT): converts $H_2O_2$ to $2H_2O$ and $O_2$ Ascorbate peroxidase (APX): converts $H_2O_2$ to $2H_2O$ Glutathione peroxidase: detoxifies $H_2O_2$ using reduced glutathione (GSH) producing the oxidised form GSSG Glutathione reductase (GR): reduces GSSG to GSH
Non enzymic	Ascorbic Acid (AA): Detoxifies $H_2O_2$ via action of APX Reduced Glutathione (GSH): co-substrate for glutathione peroxidases $\alpha$ -Tocopherol: protects against membrane lipid peroxidation (LPO) Flavonoids: scavenges $H_2O_2$ and $^1O_2$ and $OH^*$
<b>Oxidative damages to macromolecules and their cellular implications</b>	
Lipids	Oxidation of membrane lipids to products like Malondialdehyde (MDA), lipid peroxide, 4-hydroxynonenal etc) Increase in membrane fluidity and permeability
Proteins	Site-specific amino acid, modification, Fragmentation of the peptide chain, Aggregation of crosslinked, reaction products, Altered electric charge, Enzyme inactivation, Increased susceptibility, of proteins to proteolysis
DNA	Deoxyribose oxidation, Strand breakage, Removal of nucleotides, Modification of bases, DNA-protein crosslinks

**Table 3. Reports suggesting oxidative stress play a role in the phytotoxicity mechanism used by ArOPPs**

ArOPP	Plant under study	Oxidative stress/ damage markers observed	Reference(s)
Clodinafop-propargyl	Winter wheat ( <i>Triticum aestivum</i> L.); winter rye ( <i>Secale cereale</i> L., and maize ( <i>Zea mays</i> L.)	Increases in superoxide anion radical generation, LPO Increase in total antioxidant activity, catalase, and ascorbate peroxidase activity	[34]
Diclofop acid	<i>Arabidopsis thaliana</i>	High level of MDA High CAT, SOD and POD activity	[35]
Diclofop-Methyl	Rice Oat ( <i>Avena sativa</i> L.) and leafy spurge ( <i>Euphorbia esula</i> L.).	Increase in MDA, SOD and peroxidase activity Increase in MDA, SOD and peroxidase activity	[36] [37]
Fenoxaprop-ethyl	wheat	Induction of GST	[38]
Fluazifop-p (Fluazifop acid)	Sea grass ( <i>Zostera nigricaulis</i> )	Lipid peroxidation	[39]
Fluazifop-p-butyl	Bristly starbur ( <i>Acanthospermum hispidum</i> ) bristly starbur ( <i>Acanthospermum hispidum</i> ) seedlings Acanthospermum hispidum	Increase in MDA, membrane peroxidation Involvement of H <sub>2</sub> O <sub>2</sub> Membrane lipid peroxidation caused by increasing levels of reactive oxygen species (ROS)	[40] [41] [42]
Quizalofop-P-ethyl	<i>Radix isatidis</i>  <i>Lemna minor</i> and <i>Lemna gibba</i>	Reduction in SOD activity Increase in Malondialdehyde (MDA) Increase in MDA; SOD, and POD activities	[43] [44]

**Table 4. Some reports on induction of oxidative stress by ArOPPs in animal studies**

ArOPP	Animal species under study	Oxidative stress marker/ Oxidative damage observed	Reference(s)
Diclofop acid	Cyanobacteria ( <i>Microcystis aeruginosa</i> )	increase in MDA concentration and SOD activity	[47]
Fenoxaprop-ethyl	Rat	Increase in MDA and, SOD activity, decrease in GST activity, decrease in SH proteins	[49]
Fenoxaprop-p-ethyl	Cyanobacteria <i>Microcystis aeruginosa</i> and <i>Microcystis viridis</i>	generation of MDA and increase in activities of SOD, POD, CAT	[50]
Fluazifop-p-butyl	Rat	Increase in MDA level, decrease in GSH level and activities of GST, CAT, SOD	[17]
Haloxypop-P-methyl ester	Rat	Increase in MDA level, decrease in GSH level and activities of GST, CAT, SOD	[48]



resulting in lipid peroxidation and organ damage [17,48]. Table 4 summarizes the studies carried out on non-plant species linking aropps to oxidative stress.

## 5. CONCLUSION

Aryloxyphenoxypropionates are a group of very effective selective post-emergence herbicides. Their mechanism of action in target plants (inhibition of lipid synthesis and generation of oxidative stress) is unique. Presentation of similar symptoms in non-target plant and animal species will help in understanding potential development of resistance to these chemicals. These mechanisms will also contribute to effective management of clinical cases of accidental exposure to these substances.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Andrew H. Cobb, John PH. Reade herbicides and plant physiology 2<sup>nd</sup> Ed. John Wiley & Sons; 2011. ISBN 9781444322491
- Hamid AA, Aiyelaagbe OO, Balogun GA. Herbicides and its applications. *Advances in Natural and Applied Sciences*. 2011; 5(2):201-213.
- Alan wood, compendium of pesticide common names: Herbicides [Internet]; 2018. Available:[http://www.alanwood.net/pesticides/class\\_herbicides.html](http://www.alanwood.net/pesticides/class_herbicides.html) [Accessed: 2018-07-11]
- Laganà A. Herbicides (New generation): Imidazolinones, Aryloxyphenoxypropionic acids/esters and diphenylethers, 'Analysis of Pesticides'. *Encyclopedia of Analytical Chemistry: Applications, Theory and Instrumentation*. 2006;1-40. DOI: 10.1002/9780470027318.a1711
- Hamada SHE, Abdel-Lateef MF, Abdelmonem AE, El-Kholy RMA, Helalia AAR. Efficiency of certain clodinafop-propargyl formulations in controlling annual grassy weeds in wheat. *Annals of Agricultural Science*. 2013;58(1):13–18. DOI: 10.1016/j.aogas.2013.01.003
- Magani EI, Shave PA, Avav T. Evaluation of fluazifop-p-butyl and propanil for weed control in sesame (*Sesamum indicum* L.) in Southern Guinea Savanna Nigeria. *Am. J. Exp. Agric*. 2012;2(4):680–689.
- Buehring NW, Talbert RE, Baldwin FL. Rice (*Oryza sativa*) response and annual grass control with graminicides. *Weed Technol*. 2006;2:738-744. DOI: 10.1614/WT-05-153R.1.
- Lancaster ZD, Norsworthy JK, Scott RC. Evaluation of Quizalofop-Resistant Rice for Arkansas Rice Production Systems. *International Journal of Agronomy*; 2018. Article ID 6315865: 1-8. DOI: 10.1155/2018/6315865
- Nandula VK, Vencill WK. Herbicide Absorption and Translocation in Plants using Radioisotopes. *Weed Science*. 2015; 63:Special Issue:140–151. DOI: 10.1614/WS-D-13-00107.1
- Carr JE, Davies LG, Cobb AH, Pallett KE. Uptake, translocation and metabolism of fluazifop-butyl in *Setaria viridis*. *Ann. Appl. Biol*. 1986;108:15-23. DOI:10.1111/j.1744-7348.1986.tb01972.x
- Balinova AM, Lalova MP. Translocation, metabolism and residues of fluazifop-butyl in soybean plants. *Weed Research*. 1992; 32:143-147. DOI: 10.1111/j.1365-3180.1992.tb01872.x
- Ruiz-Santaella JP, Heredia A, De Prado R. Basis of selectivity of cyhalofop-butyl in *Oryza sativa* L. *Planta*. 2006;223:191–199. DOI 10.1007/s00425-005-0075-1.
- Aguero-Alvarado R, Appleby AP. Uptake, translocation and phytotoxicity of root-absorbed haloxyfop in soybean, *Festuca rubra* L. and *Festuca arundinacea* Schreb. *Weed Research* 1991;31:257-263. DOI: 10.1111/j.1365-3180.1991.tb01765.x
- Liang Y, Wang P, Liu D, Shen Z, Liu H, Jia Z, Zhou Z. Enantioselective metabolism of quizalofop-ethyl in rat. *PLoS One*. 2014 25;9(6):e101052. DOI: 10.1371/journal.pone.0101052.
- Evaluation of the new active Cyhalofop-butyl in the product barnstorm herbicide. National registration authority for agricultural and veterinary chemicals Australian Pesticides and Veterinary Medicines Authority. 2005;1-48. [ISSN1443-1335]
- Walker KA, Ridley SM, Lewis T, Harwood JL. Fluazifop a grass-selective herbicide which inhibits acetyl-coa carboxylase in sensitive plant species. *Biochemical Journal*. 1988;254(1):307–310. DOI: 10.1042/bj2551064
- Ore A, Olayinka ET Fluazifop-p-butyl, an Aryloxyphenoxypropionate herbicide, diminishes renal and hepatic functions and

- triggers testicular oxidative stress in orally exposed rats. *Toxicol Ind Health*. 2017; 33(5):406-415.  
DOI: 10.1177/0748233716657763.
18. Clark NW, Scott RC, Blain PG, Williams FM. Fate of fluazifop butyl in rat and human skin *in vitro*. *Arch Toxicol*. 1993;67(1):44-48.  
DOI: 10.1007/BF02072034
  19. Woollen BH, Hart TB, Batten PL, Laird WJ, Davies DS, Dollery CT. Oral pharmacokinetics of fluazifop-butyl in human volunteers. *Hum Exp Toxicol*. 1991; 10(1):39-43.  
DOI: 10.1177/096032719101000107
  20. Ramsey JD, Woollen BH, Auton TR, Batten PL, Leeser JE. Pharmacokinetics of fluazifop-butyl in human volunteers. II: Dermal dosing. *Hum Exp Toxicol*. 1992; 11(4):247-54.  
DOI: 10.1177/096032719201100402
  21. Secor J, Cseke C, Owen WJ. The discovery of selective inhibition of acetyl-coenzyme a carboxylase activity by two classes of graminicides. *Brighton Crop Prot. Conf. Weeds*. 1989;3B-1, 145.
  22. Rendina AR, Craig-Kennard AC, Beaudoin JD Breen MK. Inhibition of acetyl-coenzyme a carboxylase by two classes of grass-selective herbicides. *J. Agric. Food Chem*. 1990;38:1282.  
DOI: 10.1021/jf00095a029
  23. Incledon BJ, Hall CJ. Acetyl-Coenzyme A carboxylase: Quaternary structure and inhibition by graminicidal herbicides. *Pesticide Biochemistry and Physiology*. 1997;57:255-271.  
DOI:10.1006/pest.1997.2279
  24. Sasaki Y, Nagano Y. Plant acetyl-CoA carboxylase: Structure, biosynthesis, regulation and gene manipulation for plant breeding. *Biosci. Biotechnol. Biochem*. 2004;68(6):1175-1184.  
DOI: 10.1271/bbb.68.1175
  25. Heldt H. *Plant biochemistry*. 3<sup>rd</sup> Edn. Burlington, Elsevier Academic Press, USA; 2005.
  26. Lichtenthaler HK. Mode of action of herbicides affecting Acetyl-CoA carboxylase and fatty acid biosynthesis. *Biosciences*. 2014;45(5): 521-528.  
DOI: 10.1515/znc-1990-0538
  27. Hoppe HH, Zacher H. Inhibition of fatty acid biosynthesis in isolated bean and maize chloroplasts by herbicidal phenoxy-phenoxypropionic acid derivatives and structurally related compounds. *Pestic. Biochem Physiol*. 1985;24:298-305.  
DOI: 10.1016/0048-3575(85)90140-3
  28. Halliwell B. Biochemistry of oxidative stress. *Biochemical Society Transactions*. 2007;35(5):1147-1150. D  
DOI: 10.1042/BST0351147
  29. Dat J, Vandenabeele S, Vranov E, Van Montagu M, Inze D, Van Breusegem F. Dual action of the active oxygen species during plant stress responses, *Cell. Mol. Life Sci*. 2000;57:779-795.  
DOI: 10.1007/s0001800050041
  30. Waszczak C, Carmody M, Kangasjarvi J. Reactive Oxygen Species in Plant Signaling. *Annual Review of Plant Biology*. 2018;69:209-236.  
DOI:10.1146/annurev-arplant-042817-040322
  31. Demidchik V. Mechanisms of oxidative stress in plants: From classical chemistry to cell biology. *Environ. Exp. Bot*. 2014, 109:212-228.  
DOI: 10.1016/j.envexpbot.2014.06.021
  32. Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany*. 2012; 217037:1-26.  
DOI: 10.1155/2012/217037
  33. Roychoudhury A, Dasand K. Reactive Oxygen Species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Frontiers in Environmental Science: Environmental Toxicology*. 2014;2(53):1-13.  
DOI: 10.3389/fenvs.2014.00053
  34. Lukatkin AS, Gar'kova AN, Bochkarjova AS, Nushtaeva OV, Teixeira da Silva AJ Treatment with the herbicide TOPIK induces oxidative stress in cereal leaves. *Pesticide Biochemistry and Physiology*. 2013;105:44-49.  
DOI: 10.1016/j.pestbp.2012.11.006
  35. Zhang Q, Zhao M, Qian H, Lu, T, Zhang Q, Liu W. Enantioselective damage of diclofop acid mediated by oxidative stress and Acetyl-CoA carboxylase in nontarget plant *Arabidopsis thaliana*. *Environ. Sci. Technol*. 2012;46:8405-8412.  
DOI: 10.1021/es300049q
  36. Ding H, Lu H, Lavoie M, Xie J, Li Y, Lv X, Fu Z, Qian H. Unraveling the toxicity mechanisms of the herbicide diclofop-methyl in rice: Modulation of the activity of key enzymes involved in citrate

- metabolism and induction of cell membrane anion channels. *Journal of Agricultural and Food Chemistry*. 2014, 62(44):10654-10660.  
DOI: 0.1021/jf503974t
37. Shimabukuro RH, Davis DG, Hoffer BL. The effect of diclofop-methyl and its antagonist, vitamin E, on membrane lipids in oat (*Avena sativa* L.) and leafy spurge (*Euphorbia esula* L.). *Pestic. Biochem. Physiol.* 2001;69:13–26.
  38. Edwards R, Cole DJ. Glutathione transferases in wheat (*Triticum*) species with activity toward fenoxaprop-ethyl and other herbicides. *Pestic Biochem Physiol.* 1996;54:96–104.
  39. Carvea M, Coggana TL, Myersb JH, Clarkea B, Nugegodaa D, Shimeta J. Impacts on the seagrass, *Zostera nigricaulis* from the herbicide Fusilade Forte® used in the management of *Spartina anglica* infestations. *Aquatic Toxicology*. 2018;195:15–23.  
DOI:10.1016/j.aquatox.2017.11.021
  40. Luo X, Sunohara Y, Matsumoto H. Fluazifop-butyl causes membrane peroxidation in the herbicide-susceptible broad leaf weed bristly starbur (*Acanthospermum hispidum*). *Pesticide Biochemistry and Physiology*. 2004;78: 93–102.DOI:10.1016/j.pestbp.2003.10.002
  41. Luo X, Liu, Z, Sunohara, Y, Matsumoto H, Li P. Involvement of H<sub>2</sub>O<sub>2</sub> in Fluazifop-p-butyl-induced cell death in bristly starbur seedlings. *Pestic Biochem Physiol.* 2017;143:258-264.  
DOI: 10.1016/j.pestbp.2016.12.007
  42. Liu Z, Li P, Sun X, Zhou F, Yang C, Li L, Matsumoto , Luo X. Fluazifop-P-butyl induced ROS generation with IAA (indole-3-acetic acid) oxidation in *Acanthospermum hispidum* D.C. *Pestic Biochem Physiol.* 2017;143:312-318.  
DOI: 10.1016/j.pestbp.2017.10.005
  43. Xin W, Erhua R, Jinhua Z, Xiangyang Z, Yinyuan W, Pingyi G. Effect of quizalofop on protective enzymes and photosynthesis in *Radix Isatidis*. *Journal of Medicinal Plants Research* 2012;6(9):1770-1776.  
DOI: 10.5897/JMPR11.1612
  44. Doganlar ZB. Quizalofop-p-ethyl-induced phytotoxicity and genotoxicity in *Lemna minor* and *Lemna gibba*. *Journal of Environmental Science and Health, Part A*. 2012;47:1631–1643.  
DOI: 10.1080/10934529.2012.687175
  45. Tong L. Acetyl-coenzyme a carboxylase: Crucial metabolic enzyme and attractive target for drug discovery. *Cell and Molecular Biology*. 2005;62(16):1784–1803.  
DOI: 10.1007/s00018-005-5121-4
  46. Kemal C, Casida JE. Coenzyme a esters of 2-Aryloxyphenoxypropionate herbicides and 2-Arylpropionate anti-inflammatory drugs are potent and stereoselective inhibitors of rat liver Acetyl-CoA Carboxylase. *Life Science*. 1992;50(7): 533-540.
  47. Yea J, Zhang Y, Chen S, Liu C, Zhu Y, Liu W, Enantioselective changes in oxidative stress and toxin release in *Microcystis aeruginosa* exposed to chiral herbicide diclofop acid. *Aquatic Toxicology*. 2014;146:12–19.  
DOI: 10.1016/j.aquatox.2013.10.023
  48. Olayinka ET, Ore A. Hepatotoxicity, nephrotoxicity and oxidative stress in rat testis following exposure to haloxyfop-p-methyl ester, an Aryloxyphenoxypropionate herbicide. *Toxics*. 2015;3:373-389.  
DOI: 10.3390/toxics3040373.
  49. Abd-Alrahman SH, Elhalwagy MEA, Kotb GA, Farid H, Farag AAG, Draz HM, Isa AM, Sabico S. Exposure to difenoconazole, diclofop-methyl alone and combination alters oxidative stress and biochemical parameters in albino rats. *Int J Clin Exp Med*. 2014;7(10):3637-3646.
  50. Du Y, Ye1 J, Wu L, Yang C, Wang L, Hu X. Physiological effects and toxin release in *Microcystis aeruginosa* and *Microcystis viridis* exposed to herbicide fenoxaprop-p-ethyl. *Environ Sci Pollut Res*. 2017;24: 7752–7763.  
DOI: 10.1007/s11356-017-8474-y.

© 2019 Ore and Olayinka; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/48220>